

Stellar, 18.4.06  
Diamond drill hole AP004  
Collar Coordinates (GPS,AMG) 341200mE5376503mN  
RL (GPS) 119 m  
Length 152.8 m  
Azimuth (AMG) 160°  
Dip -87° degrees

Drillers' Blocks			Recovery		Geology	Foliation		Core Assays		Sample	Ni	Cu	Pb	Zn	Ag	Sn	S	Au	Pt	Element
From (m)	To (m)	(m)	From (m)	To (m)	Description	Depth	Alpha <sup>0</sup>	From (m)	To (m)	Number	ppm	ppm	ppm	ppm	ppm	ppm	%	ppm	ppm	Units
0	2.6	0.45	0.00	4.00	Clay and organics.						AAS	AAS	AAS	AAS	AAS	XRF	Leco	50 gm FA	50 gm FA	Method
2.6	4	0.00	4.00	32.40	Fine grained, schistose metasediment consisting of pale, moderately weathered, albite-muscovite-quartz-carbonate schist with metamorphic lamination comprising 1-3 mm albite-quartz bands and minus 1 mm muscovite bands. Grey due graphite in muscovite. Friable intervals due to leaching of carbonate, but pyrite is fresh. Pyrite bands up to 1 cm parallel foliation and in quartz veins at 8.8-11.8 m and 13.1-17.5 m. Intervals of dark grey richly graphitic schist (metasiltstone) at 17.9-28.5 m and 30.7-30.83 m. Core broken throughout.	4.5	25				10	10	10	10	1	10	0.01%	0.01 ppm	0.01 ppm	Sensitivity
4	5	0.30				5.5	25													
5	5.8	0.20				8.5	50													
5.8	7	0.25				19.0	45													
7	7.9	0.20				22.0	40	27	28	141132	30	120	30	290	<1	<10	1.32	<0.01	<0.01	
7.9	8.8	0.10				25.0	30	28	29	141133	30	70	20	400	<1	<10	1.58	<0.01	<0.01	
8.8	11.8	0.15				27.0	30	29	30	141134	20	20	30	370	<1	<10	0.11	<0.01	0.03	
11.8	13.1	0.00				29.5	30	30	31	141135	40	70	10	520	<1	<10	2.51	<0.01	<0.01	
13.1	17.5	0.15				32.0	15	31	32	141136	40	30	10	160	<1	<10	0.56	<0.01	0.03	
17.5	18.3	0.15	32.40	37.90	Colour becomes medium greenish-grey indicating a chloritic component in the metasediment. Minor oxidation on fractures.	33.0	22	32	33	141137	40	50	10	120	<1	<10	0.61	<0.01	0.06	
18.3	20.1	0.20				35.5	30	33	34	141138	30	30	30	130	<1	<10	0.44	<0.01	0.03	
20.1	20.4	0.30	37.90	41.80	Metasediment with narrow chloritic intervals parallel to foliation. Fresh disseminated pyrite, but minor clay and limonite on some fractures. Becomes more chloritic after 40.4 m.	37.5	40	34	35	141139	20	20	20	180	<1	<10	0.12	<0.01	0.03	
20.4	21.4	0.45						35	36	141140	30	20	20	160	<1	<10	0.14	<0.01	<0.01	
21.4	22.1	0.30						36	37	141141	30	10	20	130	<1	<10	0.07	<0.01	0.05	
22.1	23.2	0.25	41.80	46.75	Chlorite-albite-?quartz-?muscovite schist. Quartz vein boudins become prominent. Pyrite ranges from under 1% to about 5%.	44.0	30	37	38	141142	30	20	10	100	<1	<10	0.12	<0.01	<0.01	
23.2	23.7	0.25						38	39	141143	30	20	20	110	<1	<10	0.15	<0.01	0.03	
23.7	24.6	0.25	46.75	50.80	Metasediment with narrow chloritic intervals 49.8-50.8 m. Trace pyrite. Clay on fractures.	47.5	35	39	40	141144	30	20	10	160	<1	<10	0.07	<0.01	<0.01	
24.6	25.3	0.30						40	41	141145	40	30	110	340	<1	<10	1.22	<0.01	<0.01	
25.3	25.9	0.38	50.80	56.80	Chlorite-albite-?quartz-?muscovite schist. Graphite in mica generally and rich in narrow intervals. Carbonate leached and minor clay on fractures, but pyrite fresh. Core still very broken.	51.5	20	41	42	141146	30	20	10	140	<1	<10	0.76	<0.01	<0.01	
25.9	26.8	0.20				55.5	30	42	43	141147	40	30	10	220	<1	<10	1.33	<0.01	<0.01	
26.8	27.8	0.75						43	44	141148	50	50	10	130	<1	<10	0.51	<0.01	<0.01	
27.8	28.5	0.40	56.80	57.04	Milky quartz vein boudin with oxidised sulphide spots.			44	45	141149	40	30	10	330	<1	<10	0.37	<0.01	<0.01	
28.5	29.2	0.50	57.04	57.70	Albite-chlorite-?muscovite-?quartz schist. Minor pale brown carbonate veinlets near 57.7 m.	57.4	30	45	46	141150	40	80	40	310	<1	<10	4.30	<0.01	<0.01	
29.2	29.8	0.35						46	47	141151	30	80	30	280	<1	<10	3.00	<0.01	0.05	
29.8	30.7	0.50	57.70	59.70	Massive pyrite with upper contact at high angle (alpha 80) to foliation. Pyrite displays spheroidal forms with about 10% interstitial hematite and carbonate. Hematite is fine grained and may be either grey and metallic or bright cherry red and earthy. About 1% leached cavities after ?carbonate. Patchy magnetite. Chalcopyrite patches in pyrite at 59.45 m.	47	48	48	49	141152	30	30	10	210	<1	<10	0.28	<0.01	<0.01	
30.7	31.4	0.53						48	49	141153	30	30	20	140	<1	<10	0.09	<0.01	<0.01	
31.4	32.4	0.65						49	50	141154	20	20	10	90	<1	<10	0.09	<0.01	0.03	
32.4	33.5	1.05						50	51	141155	30	30	<10	130	<1	<10	0.67	<0.01	<0.01	
33.5	34.00	0.50						51	52	141156	50	40	<10	190	<1	<10	1.38	<0.01	<0.01	
34	35.1	0.85						52	53	141157	40	50	<10	160	<1	<10	1.20	<0.01	<0.01	
35.1	35.8	0.60	59.70	61.00	Sharp conformable change from random fabric to strong planar fabric due to narrow (1-3 mm) stringers of pale brown ?carbonate, fine grained magnetite, and very fine grained, white ?feldspar. Some magnetite coated by hematite, which also forms 'skins' on spheroidal pyrite. Augen textures indicate fabric is tectonic.	60.0	25	53	54	141158	40	40	<10	140	<1	<10	0.75	<0.01	0.03	
35.8	36.8	0.80						54	55	141159	30	20	<10	140	<1	<10	0.38	<0.01	<0.01	
36.8	37.9	0.75						55	56	141160	40	30	10	160	<1	<10	0.64	<0.01	<0.01	
37.9	38.8	1.10						56	57	141161	30	30	<10	130	<1	<10	0.81	<0.01	<0.01	
38.8	40.4	1.45						57	57.7	141162	30	110	<10	170	<1	<10	0.82	<0.01	<0.01	
40.4	41.8	1.40	61.00	62.30	Similar 57.7-59.7 with random to weakly oriented fabric.			57.70	58.70	140823	40	2440	110	500	6	<10	45.1	0.09	0.10	
41.8	43.7	1.85	62.30	62.60	Sharp change to planar fabric accompanied by big increase in magnetite. Increasing nonmetallic minerals downwards.			58.70	59.70	140824	40	2960	80	250	5	<10	44.5	0.07	0.02	
43.7	44.8	1.15						59.70	61.00	140825	30	2460	40	210	5	<10	32.6	0.07	0.02	
44.8	46	1.05	62.60	64.80	Dominantly massive, fine grained, quartz-feldspar rock with scattered, crude, chloritic bands that contain carbonate, pyrite and chalcopyrite.			61.00	62.00	140826	40	4260	140	240	7	<10	43.9	0.06	0.05	
46	46.9	0.85						64.0	20	140827	40	4270	90	180	5	<10	46.3	0.06	0.06	
46.9	47.8	0.55						62.30	62.60	140828	30	2600	<10	150	4	<10	21.1	0.03	0.06	
47.8	48.9	0.90	64.80	67.13	Thinly (1-10 mm) interlayered stringy bands of chloritic and felsic material with abundant pyrite and magnetite. Minor earthy red hematite, probable chalcopyrite.			62.60	63.60	140829	20	900	<10	60	3	<10	1.82	0.01	0.05	
48.9	49.8	0.90						63.60	64.65	140830	20	1800	10	100	3	<10	3.61	0.01	0.06	
49.8	50.8	0.85						64.65	65.40	140831	20	7080	<10	210	2	<10	2.63	0.01	0.07	
50.8	52.4	1.50	67.13	67.50	Pale brown carbonate vein with about 1% sulphide, mainly chalcopyrite.			65.40	66.40	140832	30	15600	190	110	4	<10	14.8	0.01	0.06	
52.4	53.8	1.40						66.40	67.13	140833	20	11800	30	190	2	<10	13.5	0.02	0.07	
53.8	54.8	0.80	67.50	68.43	Similar 64.8-67.13.			67.13	67.50	140834	30	1680	<10	100	2	<10	1.19	0.01	0.05	
54.8	55.2	0.30	68.43	70.06	Sharp change (alpha 40) to random textured, massive pyrite with minor magnetite, feldspar, chlorite and carbonate patches.			67.50	68.47	140835	30	20700	10	80	4	<10	21.3	0.02	0.07	
55.2	55.9	0.60						68.47	69.47	140836	40	11500	20	190	3	<10	34.0	0.01	0.01	
55.9	56.8	0.55	70.06	71.60	Similar 62.6-64.8.			71.0	10	140837	40	13300	<10	130	3	<10	32.9	0.03	0.10	
56.8	58.6	1.65	71.60	76.60	Breccia consisting of clasts of fine grained, white, felsic material in a massive, pyritic matrix. Clasts are up to 50 mm across and display crude planar fabric that is disoriented. Some clasts comprise thinly interlayered pyrite, hematite and magnetite. The matrix mostly has random fabric, but includes a narrow foliated band near 75.5 m. Chalcopyrite common in veinlets near 73.2 m. Bottom contact of breccia not faulted.			70.06	71.06	140838	20	18700	<10	160	2	<10	13.7	0.04	<0.01	
58.6	59.8	0.92						75.5	15	140839	20	7940	<10	160	1	<10	7.55	0.02	<0.01	
59.8	61.2	1.20						71.60	72.60	140840	30	15800	60	140	2	<10	31.3	0.03	0.04	
61.2	62.8	2.10						72.60	73.60	140841	20	13800	20	100	1	<10	15.4	0.01	0.05	
62.8	65.2	2.30						73.60	74.60	140842	20	21200	20	140	2	<10	16.5	<0.01	<0.01	
65.2	65.8	0.60						74.60	75.60	140843	30	16200	50	180	2	<10	13.3	<0.01	0.05	
65.8	68.8	2.76						75.60	76.60	140844	30	12100	60	120	2	<10	15.6	<0.01	0.03	

Drillers' Blocks	Recovery	Geology				Foliation		Core Assays		Sample	Ni	Cu	Pb	Zn	Ag	Sn	S	Au	Pt	Element	
68.8	71.8	3.20	76.60	77.00	Fine grained , chlorite-actinolite schist (?metabasalt) with disseminated magnetite and pyrite.			76.60	77.00	140845	20	7000	10	390	1	<10	2.92	<0.01	0.01		
71.8	73.2	1.50						77.00	78.00	140846	30	14800	210	290	3	<10	19.0	<0.01	<0.01		
73.2	74.8	1.56	77.00	80.80		Breccia texture with clasts up to 120 mm across. Pyritic matrix ranges from random fabric to crudely planar. Magnetite in clasts and matrix; hematite locally common; common chalcopyrite; common chlorite; common leached cavities lined with limonite.	79.1	20	78.00	79.20	140847	30	12900	110	120	4	<10	13.1	<0.01	<0.01	
74.8	77.8	2.96						79.20	80.20	140848	40	11300	<10	120	4	<10	29.5	0.02	<0.01		
77.8	80.8	3.00						80.20	80.80	140849	50	10400	<10	110	5	<10	32.2	0.01	<0.01		
80.8	83.8	2.97						80.80	81.80	140850	30	8450	<10	100	2	<10	13.7	0.01	<0.01		
83.8	86.8	2.95	80.80	85.90	Stringy chloritic, felsic and pyritic layers with massive pyrite 82.3 to 83.1. Common magnetite, pale brown carbonate. Common chalcopyrite in places. Some breccia texture.	81.0	10	81.80	82.30	140851	30	8680	<10	130	3	<10	19.4	0.01	<0.01		
86.8	89.8	2.95					81.9	20	82.30	83.10	140852	40	7940	10	120	4	<10	29.8	0.01	<0.01	
89.8	92.4	2.50					85.7	10	83.10	84.10	140853	20	7860	<10	120	2	<10	4.63	0.01	<0.01	
92.4	95.5	3.10	85.9	86.8	Probably chloritic metasandstone. Fine grained, massive , banded rock with substantial quartz, feldspar and aligned chlorite. Disseminated pyrite, but no magnetite. Common stringers of pale brown carbonate. Chalcopyrite, pyrite and pale brown carbonate present in scattered veinlets.	86.6	70	84.10	85.10	140854	40	11100	<10	90	3	<10	25.4	<0.01	<0.01		
95.5	98.6	3.10							85.10	85.90	140855	30	9940	<10	80	3	<10	23.2	<0.01	0.03	
98.6	101.6	3.00							85.90	86.90	140856	20	5740	<10	80	1	<10	5.02	0.01	<0.01	
101.6	104.7	3.16						86.90	87.86	140857	20	1230	<10	70	<1	<10	3.66	<0.01	0.02		
104.7	107.8	3.05						87.86	88.90	140858	30	1330	<10	30	<1	<10	5.07	0.02	<0.01		
107.8	109.5	1.75	86.8	95.5	Fine grained, medium grey metasandstone consisting of quartz, albite and graphitic mica with stringers of pale brown carbonate. Pyrite present, no magnetite. Minor chalcopyrite in scattered veinlets, also pale brown carbonate and quartz.	89.3	20	88.90	89.90	140859	20	1230	10	80	1	<10	2.67	0.05	0.03		
109.5	110.8	1.30					91.2	40	89.90	90.90	140860	30	1290	<10	30	1	<10	3.91	0.02	0.05	
110.8	113.8	2.80					94.4	15	90.90	91.90	140861	20	1360	<10	40	1	<10	2.91	0.01	0.05	
113.8	116.8	3.16						91.90	92.90	140862	20	1300	<10	80	1	<10	2.22	0.01	0.02		
116.8	119.8	2.70	95.5	111.55	Similar rock, but more schistose due to more abundant graphitic mica. Probably metasilstone. The mica is greenish, and may be a chlorite. Pale brown carbonate stringers persist. Thin pyrite bands present. Minor cross cutting veinlets of pyrite, quartz and carbonate. Foliation and compositional banding are discordant around 111.55 m (fold closures). Thin metasandstone intervals present throughout.	97.5	15	92.90	93.90	140863	30	1680	<10	40	<1	<10	4.05	0.01	0.03		
119.8	121.3	1.50					99.2	25	93.90	94.95	140864	30	1160	<10	40	<1	<10	3.90	<0.01	0.01	
121.3	122.8	1.50					99.9	0	94.95	95.93	140865	30	1190	<10	40	<1	<10	3.15	<0.01	0.02	
122.8	123.8	0.75					101.5	45	95.93	96.93	140866	30	720	<10	50	<1	<10	3.83	0.01	0.03	
123.8	125.1	1.30					107.7	45	96.93	98.10	140867	30	760	<10	60	<1	<10	1.99	0.01	0.05	
125.1	128.2	3.10					111.4	60	98.10	99.00	140868	50	630	10	110	1	<10	4.63	0.01	<0.01	
128.2	131.3	3.15						99.00	100.00	140869	30	440	10	110	2	<10	1.55	<0.01	0.01		
131.3	134.4	3.10	111.55	114.37	Chlorite-albite schist with rounded albite porphyroblasts up to 1 mm across. Around 5% pyrite as irregular stringy bands up to 100 mm thick in places. Pale brown carbonate occurs as stringy bands that are cut by the foliation in fold closures. Patchy magnetite and chalcopyrite present.	100.00		100.00	101.00	140870	50	490	<10	120	3	<10	3.03	<0.01	0.04		
134.4	136	1.50					101.00		102.00	140871	30	700	<10	80	2	<10	2.89	<0.01	0.04		
136	137.8	2.18					102.00		103.00	140872	30	370	<10	70	4	<10	1.77	<0.01	0.01		
137.8	140.8	3.05					103.00		104.00	140873	30	350	<10	70	3	<10	1.47	<0.01	0.04		
140.8	142.4	1.60				114.1	50	104.00	105.00	140874	40	510	<10	60	3	<10	1.55	<0.01	0.05		
142.4	143.2	0.65	114.37	117.3	Interbanded chlorite-albite schist and black, graphitic schist with a few percent pyrite in stringy bands and veinlets.	116.6	45	105.00	106.00	140875	40	600	<10	50	3	<10	3.41	<0.01	0.03		
143.2	145.3	1.80					106.00		107.05	140876	30	540	<10	50	2	<10	2.64	<0.01	0.04		
145.3	146.8	1.60	117.3	121.3	Thinly banded (1-20 mm) quartz-albite-chlorite-carbonate-pyrite schist. Common veinlets of chalcopyrite near 121.2 m.	119.7	60	107.05	108.05	140877	50	350	<10	60	2	<10	1.48	<0.01	0.03		
146.8	149.8	1.67					108.05		109.00	140878	30	620	<10	50	2	<10	2.17	<0.01	0.01		
149.8	151.9	2.05	121.3	134	Rapid, conformable transition into dolerite. Magnetite and pyrite present. Rocks become banded with intervals of chlorite schist and intervals with coarse doleritic texture preserved. Some bands albite rich. Pyrite up to 50% in places. Common magnetite. Chalcopyrite veinlets at 128.4. Common pale brown carbonate veinlets, though volumetrically minor.	121.5	45	109.00	110.00	140879	40	490	<10	40	3	<10	2.28	<0.01	0.03		
151.9	152.8	0.90					127.5	50	110.00	111.00	140880	40	420	<10	90	2	<10	3.09	0.03	<0.01	
152.8	EOH						134.0	40	111.00	111.55	140881	50	1920	<10	80	3	<10	5.04	0.02	<0.01	
							111.55		112.82	140882	40	3140	<10	140	3	<10	7.59	0.01	<0.01		
							112.82		113.62	140883	20	1130	<10	180	3	<10	1.30	0.03	0.03		
							113.62		114.37	140884	50	11300	<10	250	2	<10	13.4	0.05	0.01		
134	136.65		134	136.65	Similar chloritic schist, but more albite rich.	135.0	45	114.37	115.37	140885	20	4680	<10	70	2	<10	3.04	0.04	0.04		
136.65	137.25		136.65	137.25	Interbanded black, graphitic phyllite and massive pyrite	136.9	20	115.37	116.30	140886	40	1900	<10	70	2	<10	3.82	0.01	0.05		
137.25	137.6		137.25	137.6	Black graphitic, pyritic phyllite	137.5	55	116.30	117.30	140887	20	840	<10	130	1	<10	1.93	0.01	0.05		
137.6	137.95		137.6	137.95	Fault with puggy and coherent, black, graphitic phyllite, and breccia.	117.30		118.30	118.30	140888	40	1610	<10	170	2	<10	3.38	0.01	<0.01		
						118.30		119.30	140889	30	2030	10	130	1	<10	3.61	0.01	0.02			
137.95	148.85		137.95	148.85	Sharp conformable contact with thinly (1-10 mm) banded pale brown, fine grained quartz-carbonate rock and subordinate pale brown to cream carbonate (?dolomite). Common thin pyrite bands. Several friable, leached intervals. A little graphite in a few bands. Minor late veins of coarse grained carbonate and pyrite. Scattered chalcopyrite, but of small volume.	138.9	45	119.30	120.30	140890	30	1730	<10	60	1	<10	2.63	0.01	0.02		
						120.30		120.85	121.85	140891	30	1610	<10	50	1	<10	2.53	0.01	<0.01		
						120.85		121.85	140892	20	1020	<10	110	3	<10	2.71	0.01	0.03			
						121.85		122.88	140893	20	4030	<10	100	4	<10	4.27	0.01	<0.01			
						144.8	40	122.88	123.88	140894	40	5930	<10	260	4	<10	16.5	0.04	<0.01		
						123.88		124.90	140895	30	5940	<10	270	2	<10	10.1	0.03	0.01			
148.85	149.65		148.85	149.65	Very coarse grained, pure pyrite vein with colloform, carbonate selvages. A little quartz and red hematite in the selvages. The vein margins (alpha 10) are discordant wrt the foliation (alpha 45).	148.9	45	124.90	125.90	140896	20	3190	<10	190	2	<10	4.86	0.02	0.01		
						125.90		126.90	140897	20	3660	<10	110	4	<10	7.75	0.02	<0.01			
						126.90		127.90	140898	30	2460	<10	100	3	<10	9.82	0.02	<0.01			
149.65	152.8		149.65	152.8	Banded quartz-carbonate rock as for 137.95-148.85.	152.7	30	127.90	128.90	140899	40	2180	<10	160	4	<10	6.31	0.03	0.01		
152.8	EOH					128.90		129.90	140900	40	2750	<10	130	4	<10	5.81	0.01	0.02			
						129.90		130.90	140901	30	3470	<10	200	3	<10	5.54	0.03	<0.01			
						130.90		131.90	140902	30	2010	<10	260	3	<10	2.97	0.01	0.03			
						131.90		132.90	140903	40	4330	<10	100	4	<10	11.1	0.01	<0.01			
						132.90		133.90	140904	30	3460	<10	160	3	<10	6.82	0.02	<0.01			

Drillers' Blocks	Recovery	Geology	Foliation	Core Assays		Sample	Ni	Cu	Pb	Zn	Ag	Sn	S	Au	Pt	Element
				144.00	145.00	140955	40	2430	130	180	3	<10	11.2	0.11	0.02	
				145.00	146.00	140956	30	1950	140	170	2	<10	5.45	0.03	<0.01	
				146.00	147.00	140957	30	1910	210	200	2	<10	5.09	0.03	<0.01	
				147.00	148.00	140958	30	2030	110	150	2	<10	3.47	0.02	<0.01	
				148.00	148.80	140959	30	3570	60	100	2	<10	3.42	0.03	<0.01	
				148.80	149.65	140960	150	610	1770	120	5	<10	50.3	0.12	<0.01	
				149.65	150.60	140961	20	3720	110	190	2	<10	3.53	0.03	<0.01	
				150.60	151.60	140962	30	2650	160	190	2	<10	4.42	0.05	0.04	
				151.60	152.80	140963	30	3880	80	110	3	<10	5.77	0.04	0.04	
				152.80	EOH											

Duplicates

Sample Number	Ni ppm	Cu ppm	Pb ppm	Zn ppm	Ag ppm	Sn ppm	S %	Au ppm	Pt ppm	Element
	AAS	AAS	AAS	AAS	AAS	XRF	Leco	50 gm FA	50 gm FA	Method
	10	10	10	10	1	10	0.01%	0.01 ppm	0.01 ppm	Sensitivity
140953	20	1350	<10	80	2	n/a	n/a	n/a	n/a	
140963	n/a	n/a	n/a	n/a	n/a	<10	5.55	n/a	n/a	
140842	n/a	n/a	n/a	n/a	n/a	<10	16.2	n/a	n/a	
140843	n/a	n/a	n/a	n/a	n/a	n/a	n/a	<0.01	0.01	
140845	20	7010	<10	370	1	n/a	n/a	n/a	n/a	
140862	n/a	n/a	n/a	n/a	n/a	<10	2.16	n/a	n/a	
140864	n/a	n/a	n/a	n/a	n/a	n/a	n/a	<0.01	0.01	
140868	50	630	<10	100	1	n/a	n/a	n/a	n/a	
140882	n/a	n/a	n/a	n/a	n/a	<10	8.10	n/a	n/a	
140885	n/a	n/a	n/a	n/a	n/a	n/a	n/a	0.03	0.03	
140891	20	1600	<10	60	1	n/a	n/a	n/a	n/a	
140941	n/a	n/a	n/a	n/a	n/a	<10	2.97	n/a	n/a	
140945	n/a	n/a	n/a	n/a	n/a	n/a	n/a	0.02	<0.01	
140953	20	1350	<10	80	2	n/a	n/a	n/a	n/a	
140963	n/a	n/a	n/a	n/a	n/a	<10	5.55	n/a	n/a	
141134	n/a	n/a	n/a	n/a	n/a	n/a	n/a	<0.01	0.02	
141140	n/a	n/a	n/a	n/a	n/a	<10	0.15	n/a	n/a	
141145	40	30	90	340	<1	n/a	n/a	n/a	n/a	
141154	n/a	n/a	n/a	n/a	n/a	n/a	n/a	<0.01	0.03	
141160	n/a	n/a	n/a	n/a	n/a	<10	0.65	n/a	n/a	