


RENISON LIMITED - DRILL CORE RECORD BT 129

HOLE NUMBER	BT 129	SURVEY			From - To	Distance D	VERTICAL		HORIZONTAL	
		Depth	Bearing	Dip			D. Sin. Dip	H.L.	D. Cos. Dip	Prog. Total
PURPOSE	To test for extensions of Anchor mineralization		GRID					256.44		
		0	Casing	- 56°	0 - 10	10	8.290	248.15	5.592	5.59
		41m	321	- 56.25	10 - 62	52	43.236	204.91	28.890	34.48
		83m	323	- 56.5	62 - 83	21	17.512	187.40	11.591	46.07
LOCATION	Anchor Open Cut									
COLLAR R.L.	256.44									
CO-ORDINATES	5240.75mN 4675.87mE									
LENGTH	83m									
HOLE SIZE	0 - 6m NQ -83m BQ									
DATE DRILLED	7.4.81 to 8.4.81									
SIGNIFICANT CORE LOSS ZONES										
ORE ZONE GROUND CONDITIONS										
LOGGED BY	A. ROSS									
COMMENTS	Interval from 0 to 83m assayed. Several zones of very low grade Sn were encountered.									

SUMMARY - ASSAY DATA


LODE NAME	FROM	TO	LENGTH (m)	AVERAGE WEIGHTED ASSAYS										B.C.A.	
				Sn.	Acid Sol. Sn.	Cu.	As.	S.	Pb.	Zn.	Bi.	WO <sub>3</sub>	Ag g/t		
0.1% Cut-off	0	2	2	0.16		< 0.01					< 0.01		< 1		
	(256.4RL)	(254.8RL)	(1.6 BTT)												
0.1% Cut-off	6	13	7	0.21		< 0.01					0.02		< 1		
	(251.5RL)	(245.7RL)	(5.8BTT)												
0.1% Cut-off	17	22	5	0.31		< 0.01					< 0.01		< 1		
	(242.3RL)	(238.2RL)	(4.1BTT)												
0.1% Cut-off	26	33	7	0.20		< 0.01					< 0.01		< 1		
	(234.8RL)	(229.0RL)	(5.8 BTT)												

HOLE No.: BT 129




RENISON LIMITED  
DIAMOND DRILL HOLE PLOT

BT 129




PLAN


5277.0 N  
4647.4 E



5240.75 N  
4675.87 E



DIP PROFILE



892117

## DIAMOND DRILL RECORD

HOLE NUMBER : BT 129

LOGGED BY : AFR

NWFS

INTERVAL (m)		RECOVERY		DESCRIPTION	FORM	% Sn.										
FROM	TO	m	%			FROM	TO	TOTAL	ACID SOL.	% Cu.	% As.	% Mn	% Pb.	% Zn.	% Bi.	g/t Ag
				SUMMARISED LOG		0	1	0.19	0.0035		0.035		0.0060		1	
							2	0.13	0.0045		0.035		0.0090		<1	
0	69.7			VARIABLY ALTERED ALKALI GRANITE WITH ZONES OF GRANITE-GREISEN, GRANULAR GREISEN, FELDSPATHISED GRANITE (ANCHOR GRANITE).			3	0.05	0.0120		0.04		0.0045		2	
							4	0.03	0.0055		0.04		0.0050		1	
							5	"	0.0040		0.035		0.0035		1	
69.7	83.0			WEAKLY ALTERED, GREISEN-GRANITE TO GRANITE-GREISEN.			6	0.02	0.0055		0.03		0.0050		1	
							7	0.23	0.0045		0.03		0.0060		1	
							8	0.33	0.0020		0.03		0.0105		1	
				DETAILED LOG			9	0.34	0.0015		0.04		0.0140		<1	
							10	0.13	0.0015		0.035		0.0120		<1	
0	1.0	1.0	100	Broken, limonitic weathered joints. Grey cream equigranular, medium grained greisen alkali granite. No obvious cassiterite.			11	0.12	0.0015		0.035		0.0135		<1	
							12	0.17	0.0015		0.04		0.0075		<1	
							13	0.16	0.0010		0.05		0.0070		<1	
1.0	5.8	4.8	100	Clayey joints. Grey cream greisen granite. No obvious cassiterite. Sparse disseminated micas.			14	0.08	0.0010		0.045		0.0055		<1	
							15	0.03	0.0010		0.05		0.0070		<1	
							16	0.02	0.0010		0.045		0.0060		<1	
5.8	6.4	0.6	100	Whiter cream granite-greisen.			17	0.01	0.0010		0.04		0.0050		<1	
							18	0.11	0.0010		0.045		0.0060		<1	
6.4	9.5	3.1	100	Alternating mixture of grey cream granite-greisen and greyer greisen- granite. No obvious cassiterite.			19	0.56	0.0010		0.045		0.0060		<1	
							20	0.60	0.0010		0.04		0.0050		<1	
							21	0.12	0.0010		0.035		0.0060		<1	
9.5	16.9	7.4	100	Grades into grey yellow greisen granite with irregular patches of alteration, similar to BT 57, e.g. fluorite/mica patches to 5cm. Minor clayey joints.			22	0.16	0.0015		0.045		0.0060		<1	
							23	0.02	0.0015		0.05		0.0080		<1	
							24	<0.01	0.0010		0.045		0.0070		<1	
							25	0.01	0.0010		0.05		0.0105		<1	
16.9	21.1	4.2	100	Grey cream sericitised granite greisen. Less altered than the previous zone.			26	0.05	0.0015		0.04		0.0060		<1	
							27	0.26	0.0015		0.025		0.0060		<1	
							28	0.14	0.0010		0.05		0.0075		<1	
21.1	24.7	3.6	100	Grades into zone of more intense alteration. Grey-green to pale grey granular greisen to greisen-granite. No obvious cassiterite but could be expected to occur in this rock type.			29	0.03	0.0015		0.06		0.0070		<1	
							30	0.12	0.0015		0.035		0.0060		<1	
							31	0.26	0.0015		0.035		0.0060		<1	
							32	0.25	0.0010		0.035		0.0045		<1	
24.7	27.5	2.8	100	Grades to lesser altered, grey cream granite-greisen.			33	0.37	0.0015		0.035		0.0045		<1	
							34	0.06	0.0010		0.03		0.0045		<1	
27.5	28.9	1.4	100	Grades to grey, more altered pale grey granular greisen with coarse micas. No obvious cassiterite.			35	<0.01	0.0010		0.05		0.0045		<1	
							36	"	0.0010		0.025		0.0050		<1	
							37	"	0.0010		0.03		0.0060		<1	
28.9	33.7	3.8	100	Grades to grey-cream granite-greisen with sparse coarse micas. No obvious cassiterite.			38	"	0.0010		0.045		0.0060		<1	
							39	"	0.0010		0.065		0.0085		<1	
							40	"	0.0010		0.04		0.0045		<1	
33.7	34.2	0.5	100	Variable zone of slight pink granite greisen to greyish more feldspathic granite-greisen.			41	"								
							42	"								

892118

## DIAMOND DRILL RECORD

HOLE NUMBER : BT 129

LOGGED BY : AFR

NWPS

INTERVAL (m)		RECOVERY		DESCRIPTION	FORM.	% Sn.										
FROM	TO	m	%			FROM	TO	TOTAL	ACID SOL.	% Cu.	% As.	% Mn	% Pb.	% Zn.	% Bi.	g/t Ag
34.2	34.5	0.3	100	Zone of intense alteration. Grey granular greisen with a basal "contact" marked by abundant mica at 55° CA.		42	43	< 0.01								
							44	"								
							45	"								
34.5	36.9	2.4	100	Intense zone of feldspar rich altered granite. Pale pink to lime green feldspar-greisen-granite with sparse dark mica.			46	"								
							47	"								
							48	"								
36.9	37.1	0.2	100	More normal but pinkish greisen-granite to granite greisen. Medium grained equigranular.			49	"								
							50	"								
							51	"								
37.1	39.25	2.15	100	Intense feldspar zone. Pinkish feldspar rich granite with sparse disseminated mica. No obvious cassiterite.			52	"								
							53	"								
							54	"								
39.25	46.0	6.75	100	Grades into greyish green, variable greisen-granite to granite greisen. Grey green to cream-grey in colour. Minor zones approaching granular greisen texture. Pinkish in parts.			55	"								
							56	"								
							57	"								
							58	"								
46.0	53.3	7.3	100	Colour becomes more greyish-green. Variable greisen granite to almost granular greisen. Siderite common.			59	0.01								
							60	< 0.01								
							61	"								
53.3	58.1	4.8	100	Pronounced zone of pinkish cream feldspar rich granite with sparse micas, with minor zone of greenish sericitised greisen-granite.			62	"								
							63	"								
							64	"								
58.1	59.1	1.0	100	Zone of intense mica greisen. Dark grey green. Contacts gradual but "sharp".			65	"								
							66	"								
							67	"								
59.1	67.0	7.9	100	Grades into variable pinkish granite greisen to grey green granular greisen (granite) with coarse dark feldspars. Siderite present in parts. From 63m the rock is more of a greisen granite. Minor clayey zones.			68	"								
							69	"								
							70	"								
							71	"								
							72	"								
67.0	69.2	2.2	100	Grades to a more pink granite - weak greisen.			73	"								
							74	"								
69.2	69.7	0.5	100	Grades into a more grey green greisen-granite to granular greisen. Gradational contacts.			75	"								
							76	"								
							77	"								
69.7	83.0	13.3	100	Variable but mainly greyish greisen-granite with minor zones of darker grey-green greisen-granite, and zones of pinkish-cream feldspar rich granite (greisen). Minor siderite. Speck moly.			78	"								
							79	"								
							80	"								
							81	"								
				END OF HOLE			82	"								
							83	"								

Sn Assays by Mines Dept., Lunenburg (XRF)  
 Cu, Zn, Ag, Mn Assays by Renison (AAE)

892119