


RENISON LIMITED - DRILL CORE RECORD

| HOLE NUMBER | BT 80 | SURVEY | | | From - To | Distance D | VERTICAL | | HORIZONTAL | |
|-----------------------------|--|--------|------------|-------|-----------|------------|-----------|-------|------------|-------------|
| | | Depth | Bearing | Dip | | | D.Sin.Dip | R.L. | D.Cos.Dip | Prog. Total |
| PURPOSE | To test any southern extension of mineralization. | 40m | 016.5° mag | 87.5° | 0 | | | 280.3 | | |
| | | 82m | 015° mag | 88° | 83 | | | 197.3 | | |
| LOCATION | Anchor Mine | | | | | | | | | |
| COLIAR R.L. | 280.3m | | | | | | | | | |
| CO-ORDINATES | 5249.7m N; 5014.6m E | | | | | | | | | |
| LENGTH | 83m | | | | | | | | | |
| HOLE SIZE | 0 - 18m N.Q. CORE 18m - 83m B.Q. CORE | | | | | | | | | |
| DATE DRILLED | 24.4.80 - 1.5.80 | | | | | | | | | |
| SIGNIFICANT CORE LOSS ZONES | | | | | | | | | | |
| ORE ZONE GROUND CONDITIONS | | | | | | | | | | |
| LOGGED BY | L. MARTIN / | | | | | | | | | |
| COMMENTS | <p><i>Very low grade tin zones encountered.</i></p> <p><i>The shallower of these contains high Ag values. (AFR)</i></p> <p><i>12m NQ casing left (AFR)</i></p> | | | | | | | | | |

SUMMARY - ASSAY DATA

| LODE NAME | FROM | TO | LENGTH (m) | AVERAGE WEIGHTED ASSAYS | | | | | | | | | | | B.C.A. |
|--------------|------|------|------------|-------------------------|---------------|------|-------|------|-------|-------|-------|-----------------|--------|-------|--------|
| | | | | Sn. % | Acid Sol. Sn. | Cu. | As. | S. | Pb. | Zn. | Bi. | WO ₃ | Ag g/t | Mo | |
| 0.1% Cut Off | 33.4 | 38.4 | 5 | 0.28 | 0.01 | 0.29 | <0.10 | 0.1 | <0.01 | 0.02 | 0.005 | <0.01 | 32 | 0.002 | |
| | 53.5 | 59.5 | 6 | 0.12 | 0.01 | 0.02 | <0.1 | <0.1 | <0.01 | <0.01 | 0.002 | <0.01 | 1 | 0.002 | |
| | | | | | | | | | | | | | | | |
| | | | | | | | | | | | | | | | |


FILE No. BT80



REINSON LIMITED DIAMOND DRILL HOLE PLOT


5 cm

PLAN



DIP PROFILE

380.3 m



0.00
1.00
2.00
3.00
4.00
5.00
6.00
7.00
8.00
9.00

10.00
11.00
12.00
13.00
14.00
15.00
16.00
17.00
18.00
19.00
20.00
21.00
22.00
23.00
24.00
25.00
26.00
27.00
28.00
29.00
30.00
31.00
32.00
33.00
34.00
35.00
36.00
37.00
38.00
39.00
40.00

197.3 m

41.00
42.00
43.00
44.00
45.00
46.00
47.00
48.00
49.00
50.00
51.00
52.00
53.00
54.00
55.00
56.00
57.00
58.00
59.00
60.00
61.00
62.00
63.00
64.00
65.00
66.00
67.00
68.00
69.00
70.00
71.00
72.00
73.00
74.00
75.00
76.00
77.00
78.00
79.00
80.00
81.00
82.00
83.00
84.00
85.00
86.00
87.00
88.00
89.00
90.00
91.00
92.00
93.00
94.00
95.00
96.00
97.00
98.00
99.00
100.00

038027

DIAMOND DRILL RECORD

HOLE NUMBER : BT 60

LOGGED BY : L.MARTIN

NWPS

| INTERVAL (m) | | RECOVERY | | DESCRIPTION | FORM | % Sn | | | | | | | | | | |
|--------------|-------|----------|-----|--|------|------|------|-------|-----------|-------|-------|-------|-------|--------|-------|--------|
| FROM | TO | m | % | | | FROM | TO | TOTAL | ACID SOL. | % Cu. | % As. | % S. | % Pb. | % Zn. | % Bi. | g/t Ag |
| 0 | 15.4 | - | - | SAND Coarse grained, orange brown, iron oxide stained. Minor weathered Poimena granite rubble. | | | | | | | | | | | | |
| 15.4 | 32.48 | | | POIMENA GRANITE: | | | | | | | | | | | | |
| | | 0.14 | 95 | 15.4 - 15.55m : Coarse grained porphyritic, orange brown weathered and iron-oxide stained; some fracturing at 45°. | | 26 | 33 | <0.10 | | | | | | | | |
| | | 3.51 | 95 | 15.55 - 19.25m : Fresh, grey and hard. Bond of greenish muscovite at 70°, 3cm wide at 17.35m. Fractured zone at 17.4-17.6m with altered yellow chlorite fillings. | | *33 | 39 | >0.10 | | | | | | | | |
| | | 8.12 | 95 | 19.25 - 27.8m : Light pinkish grey rock, very fractured and crumbly with feldspars altered to light green clays. Aplite vein at 27.36 - 27.48m, composed of fine grained quartz and feldspars, with gradational boundaries to Poimena granite. | | 39 | 49 | <0.10 | | | | | | | | |
| | | 3.52 | 98 | 27.8m - 31.40m : Pink and slightly altered with minor green chlorite filled fractures. Pegmatite vein at 27.9 - 28.18m, composed of coarse grained quartz, muscovite, black biotite and minor feldspar; sub-horizontal boundaries. Aplite zones at 29.1 - 29.47m and 30.39 - 30.94m, fine grained grey-green, with vague bands slightly richer in biotite. Diffuse boundaries over several cm. | | *49 | 50 | 0.10 | | | | | | | | |
| | | 1.08 | 100 | 31.40 - 32.48m : Grey and fresh. | | 50 | 54 | <0.10 | | | | | | | | |
| 32.48 | 32.62 | 0.14 | 100 | PEGMATITE: Medium grained pinkish feldspar, quartz and asicular biotite and muscovite. Vague pink and grey banding; gradational boundaries over 2cm and sub horizontal. Fractured core with light green chlorite and talc fillings. | | *63 | 67 | >0.10 | | | | | | | | |
| 32.62 | 33.0 | 0.38 | 100 | APLITE: Cream, fine grained quartz feldspar rock with minor sub horizontal banding; gradational boundaries. | | 67 | 93 | <0.10 | | | | | | | | |
| 33.0 | 41.78 | 8.78 | 100 | LOPTAH GRANITE - GREISEN: 33.0 - 38.5m : medium to coarse grained, pink grading to grey zones where BORTITE and coarse grained biotite occur interstitially. Minor green chlorite talc filled fractures at 40° - 50°. (Pit 33.90-33.96m) 38.5 - 41.78m : Medium to coarse grained, slightly altered with pink feldspars. Purple fluozite - chlorite vein at 41.7m, 30° with slicken-sides. | | 32.4 | 33.4 | 0.09 | <0.1 | <0.1 | <0.01 | <0.01 | | | | |
| | | | | | | 34.4 | 35.4 | 0.07 | <0.1 | <0.1 | <0.01 | 0.01 | | See p3 | <0.01 | |
| | | | | | | 35.4 | 36.4 | 0.02 | <0.1 | <0.1 | <0.01 | 0.01 | | | <0.01 | |
| | | | | | | 36.4 | 37.4 | 0.09 | <0.1 | 0.2 | <0.01 | 0.03 | | | <0.01 | |
| | | | | | | 37.4 | 38.4 | 0.20 | <0.1 | <0.1 | <0.01 | 0.02 | | | <0.01 | |
| | | | | | | 38.4 | 39.4 | 0.02 | <0.1 | <0.1 | <0.01 | 0.01 | | | <0.01 | |
| | | | | | | 39.4 | 42.6 | <0.01 | <0.1 | <0.1 | <0.01 | <0.01 | | | <0.01 | |

038025

DIAMOND DRILL RECORD

HOLE NUMBER : BT 60

LOGGED BY : L.MARTIN

KWPS

| INTERVAL (m) | | RECOVERY | | DESCRIPTION | FORM. | % Sn. | | | | | | | | | | | |
|--------------|-------|----------|-----|--|-------|-------|-------|-------|----------|-------|-------|-------|-------|-------|-------|--------|-------------------|
| FROM | TO | m | % | | | FROM | TO | TOTAL | ACID SOL | % Cu. | % As. | % S. | % Pb. | % Zn. | % Bi. | g/t Ag | % WO ₃ |
| 41.78 | 46.98 | 5.2 | 100 | <u>POIMENA GRANITE:</u> 41.78 - 43.22m : Coarse grained, porphyritic with salmon pink feldspars. Upper contact is sub horizontal with biotite enrichment for about 8 cm in Poimena granite. Mixed pegmatite and aplite zone at 42.06 - 42.42m, with BORNITE, chlorite, and fluorite. Boundaries are sub-horizontal. 43.22 - 46.98m: Fresh and grey. Pinkish and crumbly for last 20cm. | | | | | | | | | | | | | |
| 46.98 | 47.51 | 0.53 | 100 | <u>PEGMATITE:</u> Very coarse grained salmon pink feldspar and creamy quartz to 10cm diameter. Minor irregular zones of finer grained pegmatite richer in asicular black biotite. | 46.5 | 49.5 | <0.01 | | <0.1 | <0.1 | <0.01 | <0.01 | | | | <0.01 | |
| 47.51 | 53.95 | 6.44 | 100 | <u>APLITE/FINE GRAINED GRANITE:</u> 47.51 - 49.65m: Very fine grained, greenish, altered, crumbly, and fractured rock with large angular fragments of pink feldspar mixed in. 49.65 - 49.78m: Dark green biotite rock with gradational boundaries. 49.78 - 53.95m: Fine grained and greenish, grading into fine grained creamy quartz - feldspar rock with light grey diffuse patches and minor dark biotite spots. Gradational lower boundary with biotite enrichment for several cm. | 49.5 | 50.5 | 0.02 | | <0.1 | <0.1 | <0.01 | <0.01 | | | | <0.01 | |
| | | | | | 50.5 | 51.5 | 0.01 | | <0.1 | <0.1 | <0.01 | <0.01 | | | | <0.01 | |
| | | | | | 51.5 | 53.5 | <0.01 | | <0.1 | <0.1 | <0.01 | <0.01 | | | | <0.01 | |
| 53.95 | 59.6 | 5.65 | 100 | <u>LOTTAH GRANITE:</u> Fine grained, gray to light brown, fresh and hard; minor fractures. (P& 59.01 - 59.06m) | 53.5 | 54.5 | 0.12 | | <0.1 | <0.1 | <0.01 | <0.01 | | | | <0.01 | |
| | | | | | 54.5 | 55.5 | 0.06 | | <0.1 | <0.1 | <0.01 | <0.01 | | | | <0.01 | |
| | | | | | 55.5 | 56.5 | 0.01 | | <0.1 | <0.1 | <0.01 | <0.01 | | | | <0.01 | |
| 59.6 | 61.18 | 1.58 | 100 | <u>QUARTZ- FELDSPAR ROCK:</u> Fine grained, cream with grey irregular patches and biotite spots. Crumbly core with green muscovite rich zones; gradational boundaries. | 56.5 | 57.5 | 0.03 | | <0.1 | <0.1 | <0.01 | <0.01 | | | | <0.01 | |
| | | | | | 57.5 | 58.5 | 0.18 | | <0.1 | <0.1 | <0.01 | <0.01 | | | | <0.01 | |
| | | | | | 58.5 | 59.5 | 0.30 | | <0.1 | <0.1 | <0.01 | <0.01 | | | | <0.01 | |
| | | | | | 59.5 | 60.5 | 0.02 | | <0.1 | <0.1 | <0.01 | <0.01 | | | | <0.01 | |
| | | | | | 60.5 | 61.5 | 0.01 | | <0.1 | <0.1 | <0.01 | <0.01 | | | | <0.01 | |
| 61.18 | 83.0 | 21.82 | 100 | <u>LOTTAH GRANITE:</u> 61.18 - 64.5m : Fine grained, grey to light brown, fresh and hard. 64.5 - 65.15m : Light green, altered and crumbly. 65.15 - 66.96m : As for 61.18 - 64.5m 66.96 - 67.35m : Dark pink feldspars becoming pinker around a pegmatite - clay - chlorite vein at 67.18m, 1cm wide at 70°. 67.35 - 83.0m : Medium grained light brown to grey, fresh and hard. Minor fractures at 30°-50° with yellow altered chlorite fillings. Biotite zone at 71.18 - 71.55, 2cm wide, at 10°. Greisen veins at 78.75 - 78.8 and 81.6 - 82.44, 3-4cm wide at 40°-50°, fine grained, grey and sugary textured. | 61.5 | 62.5 | 0.03 | | <0.1 | <0.1 | <0.01 | <0.01 | | | | | <0.01 |
| | | | | | 62.5 | 63.5 | 0.02 | | <0.1 | <0.1 | <0.01 | <0.01 | | | | <0.01 | |
| | | | | | 63.5 | 64.5 | 0.08 | | <0.1 | <0.1 | <0.01 | <0.01 | | | | <0.01 | |
| | | | | | 64.5 | 65.5 | 0.05 | | <0.1 | <0.1 | <0.01 | <0.01 | | | | <0.01 | |
| | | | | | 65.5 | 66.5 | 0.10 | | <0.1 | <0.1 | <0.01 | <0.01 | | | | <0.01 | |
| | | | | | 66.5 | 67.5 | 0.03 | | <0.1 | <0.1 | <0.01 | <0.01 | | | | <0.01 | |
| | | | | | 67.5 | 68.5 | 0.01 | | <0.1 | <0.1 | <0.01 | <0.01 | | | | <0.01 | |
| | | | | | 68.5 | 69.5 | <0.01 | | <0.1 | <0.1 | <0.01 | <0.01 | | | | <0.01 | |
| | | | | | 81.0 | 83.0 | <0.01 | | <0.1 | <0.1 | <0.01 | <0.01 | | | | <0.01 | |

END OF HOLE

038029

