

LOCATION	Sterling Valley	Depth(m)	Direction	Dip.	Depth(m)	Direction	Dip.	COLLAR DIP. -60°	TOTAL DEPTH 249.1m
OBJECTIVE	To test a coincident magnetic & I.P. anomaly beneath glacial overburden	30	-	-63°	200	109°	-40°	DIRECTION 108°	HOLE SIZE NQ 48m BQ 249.1m
RESULT	Tin assays: 92.35m - 93.35m, 4950 ppm; 98.65-105.45 averaged 1400 ppm	66	109°	-60°	249	108°	-26.5°	R.L. 176m	COMMENCED 3.4.'80
		84	112°	-58.5°				COORDINATES	COMPLETED 2.5.'80
		120	112°	-56°				4280N 4565E S.V.G.	LOGGED BY A.J. Mollison
		150	108°	-50°				384,190mE 5,374,393mN	

DEPTH (m)		ROCK DESCRIPTION REFER DDH SUMMARY SHEET A1-526-0012	MINERALISATION	SAMPLE NO.	FROM	TO	CORE REC'D	ASSAY DATA per ppm							CORE REC'D		
FROM	TO							Sample Length	Pb.	Zn.	Cu	Ag - g/t	As	Fe.	Sn.	RUN	SHORT
0	17.2	GLACIAL OVERBURDEN	Sulphides negligible. Trace fresh pyrite at 47.4m and 57.1m												0-1.5	0.5	
17.2	58.9	TRACHY-ANDESITE dark green fine grained, probably intrusive, unit, which is magnetic throughout.	Remnant sulphide textures in oxidised core												3	1.0	
17.2	36.4	Heavily weathered section of unit. Orange-brown with remnant feldsphyric texture locally epidotised and epidote veins. Joints strongly limonitic and manganiferous.													4	0.2	
36.4	45	As above - less weathered becoming fresher down hole, with a blochy yellow green colour becoming more uniformly green. Cut locally by remnant quartz veins now filled limonite. The texture of these veins suggests the limonite may be after sulphides.													4.5	0.4	
45	58.9	As above - dark green, fine grained porphyritic in places, with tension fractures filled by epidote with later quartz, carbonate (calcite and rhodocrosite), chlorite and fluorite alteration.													5.5	0.4	
58.9	74.5	MICRO GABBRO dark green medium grained holc crystalline unit extensively cut by veins of quartz carbonate, chlorite, and possibly fluorite and hematite													11.6	6.0	
70.4	73.2	Shear zone Micro Gabbro heavily cleaved													14.6	3.8	
74.5	98.5	ANDESITE green grey fine grained lava													16	1.7	
74.5	79	As above with frequent carbonate, quartz and chlorite veining, and associated shearing.													16.6	0.4	
				33081	0	5	c	5.0	10	30	25	x	8		A9	17.2	0.6
				082	5	10	c	5.0	15	30	5	x	10		A1	28.6	9.0
				083	10	15	c	5.0	10	25	5	1.0	16		A9	30.1	0.3
				084	15	20	c	5.0	50	165	30	x	46		A1	31.6	0.5
				085	20	25	c	5.0	15	125	25	x	10		A9	33.1	0.4
				086	25	30	c	5.0	10	140	115	x	7		A2	36.1	1.5
				087	30	35	c	5.0	10	105	80	x	10		A1	39.1	1.3
				088	35	40	c	5.0	5	140	80	0.5	4		A1	40	0.2
				089	40	45	c	5.0	15	125	50	1.0	15		A2	40.5	0.1
				33090	45	50	c	5.0	10	135	200	x	7		A4	42.5	0.1
				091	50	55	c	5.0	15	115	145	x	6		A2	45.5	0.3
				092	55	60	c	5.0	10	180	305	x	6		A5	46.1	-
																48.1	-
																51.1	-
				29797	51.10m											54.1	-
																57.1	-
				37954	60	61	sc	1.0	5	145	140	x	10	0.02	A2	58.9	-
				955	61	62	sc	1.0	x	150	60	1.0	10	0.01	A2	63.1	0.6
				956	62	63	sc	1.0	x	150	35	1.0	13		F490	66.1	-
				957	63	64	sc	1.0	x	140	55	x	3		A3	69.1	-
				958	64	65	sc	1.0	20	130	105	0.5	4		A4	70.3	-
				959	65	66	sc	1.0	10	105	220	x	4		A6	71.1	0.3
				37960	66	67	sc	1.0	10	160	110	x	2		A5	72.1	0.6
				961	67	68	sc	1.0	15	215	65	x	4		A8	73.2	0.5
				962	68	69	sc	1.0	20	140	160	x	8		A4	75.1	-
				963	69	70	sc	1.0	20	105	10	x	8		A4	78.1	0.2
				964	70	71	sc	1.0	40	175	1600	2.5	620		F2000	81.1	0.2
				965	71	72	sc	1.0	15	160	30	x	20		A10	84.1	-
				966	72	73	sc	1.0	15	165	70	x	21		A11	87.1	-
				967	73	74	sc	1.0	10	215	40	0.5	44		A4	89.5	-
				968	74	75	sc	1.0	25	170	80	1.0	4		A6		-
				969	75	76	sc	1.0	50	180	150	1.0	4000		F860		-

DEPTH (m)		ROCK DESCRIPTION	MINERALISATION	SAMPLE NO.	FROM	TO	CORE REC'D	ASSAY DATA							CORE REC'D		
FROM	TO							Sample Length	Pb	Zn	Cu	Ag - g/t	per As	ppm Fe	Sn	RUN	SHORT
79	82.1	As above with minor carbonate veins (less than 5% of rock) with 10 cm wide band of strong silica, chlorite alteration at 79.5m	<u>79.0-82.1</u> Approximately 0.1% pyrite as cubic infilling up to 5mm diam	37970	76	77	sc	1.0	10	155	50	x	24	Au	F45	90.1	-
				971	77	78	sc	1.0	30	180	90	0.5	24	As	F270	91.1	0.1
				972	78	79	sc	1.0	20	100	155	1.5	4000	3000	F1000	96.1	-
				973	79	80	sc	1.0	80	600	310	2.5	70		Fx	99.1	-
82.1	88.2	As above with minor fine carbonate quartz and epidote veins forming approx 2% of rock. <i>87-1 T/S fg dioritic intrusion (2.6m)</i>	<u>82.1-88.2</u> Trace pyrite & pyrrhotite	974	80	81	sc	1.0	35	185	100	1.3	12		F25	100.6	-
				975	81	82	sc	1.0	30	200	40	1.5	14		F270	102.1	-
				976	82	83	sc	1.0	10	110	15	1.0	4		Fx	105.1	-
88.2	89.5	As above more frequent carbonate & quartz veins up to 20% of rock averaging 5% with veins up to 1cm wide	<u>88.2-89.5</u> Pyrite vein infillings and cubes up to 5mm diam up to 0.5% volume with associated pyrrhotite (0.1% volume).	977	83	84	sc	1.0	20	135	25	1.5	10		F80	108.1	-
				978	84	85	sc	1.0	50	230	45	1.0	8		F310	111.1	-
				979	85	86	sc	1.0	40	200	180	x	22		F630	114.1	-
				37980	86	87	sc	1.0	20	180	100	0.5	12		F330	117.1	-
				981	87	88	sc	1.0	20	135	15	x	5		Ax	120.1	-
				982	88	89	sc	1.0	15	110	20	x	4		Ax	123.1	-
74.5	98.5	ANDESITE green grey fine grained lava														126.1	-
89.5	92.3	As above - heavily cleaved, chloritised, sericitised and otherwise altered rock. Feldspar grains show alignment parallel to cleavage.	Pyrite & pyrrhotite occur in approximately equal proportions up to 50% volume in veins up to 3cm wide with an average of 5% total pyrite & pyrrhotite.	29798	87.10		Thin	Section - orthoclase microdiorite							129.1	0.1	
				33099	89	90	sc	1.0	25	185	420	x	580		A5	138.1	-
				33100	90	90.65	sc	0.65	20	230	70	1.0	8		A1	138.7	-
				33066	90.65	92.35	sc x	0.7	5	130	470	0.5	136	0.18	Fx	141.1	-
92.3	98.5	Common (approximately 5% of rock) narrow carbonate, quartz veins (up to 1cm wide) in a weakly but pervasively altered andesite <i>93.1 - T/S. Breccia - Tourmaline, quartz, arsenopyrite</i>	Pyrrhotite with pyrite and minor chalcopyrite. Pyrrhotite veins average 5% volume of rock & are up to 1cm wide pyrite occasionally associated with pyrrhotite with rare chalcopyrite	067	92.35	93.35	sc x	1.0	535	440	1550	3.5	1.8%	0.21	F4950	144.1	-
				068	93.35	94.35	sc x	1.0	35	165	900	1.0	305	x	F60	147.1	-
				069	94.35	95.35	sc x	1.0	5	200	325	0.5	160	x	F170	150.1	-
				33070	95.35	96.36	sc x	1.0	10	210	400	x	112	x	F60	153.1	-
				071	96.35	97.35	sc x	1.0	15	155	350	x	76	x	F30	156.1	-
				072	97.35	98.65	sc x	0.9/1.3	5	155	380	0.5	920	0.01	F360	162.1	-
				073	98.65	99.65	sc x	1.0	40	115	2360	4.5	6%	0.91	F3700	163.1	-
98.5	100.6	Massive arsenopyrite and pyrrhotite with minor pyrite, chalcopyrite and cassiterite in a silica-chlorite-carbonate-fluorite gangue <i>100.2 T/S Semi-massive sulphide rock</i> <i>100.65 T/S qb - tourmaline - sulphide rock</i>	Up to 100% total sulphides with approximately 40% arsenopyrite 40% pyrrhotite 15% pyrite & 5% chalcopyrite with traces of cassiterite. Arsenopyrite & pyrrhotite occur as broad stringers up to 0.5m wide with pyrite fringing the stringers. Chalcopyrite is associated with pyrrhotite and pyrite as small grains and trains of grains up to 2mm diameter. Cassiterite occur as very fine grains at the interface between the gangue minerals and the major mineralisation.	074	99.65	100.7	sc x	1.05	140	125	2250	4.5	15%	*4.01	F1550	165.1	-
																168.1	-
																171.1	-
																174.1	-
																177.1	-
																180.1	0.1
																183.1	0.3
				33075	100.7	101.65	sc x	0.95	25	170	465	1.5	2700	0.09	F360	186.2	0.1
				076	101.65	102.55	sc x	0.9	20	1050	1200	3.0	3.5%	0.25	F1500	189.1	0.2
				077	102.55	103.55	sc x	1.0	5	110	285	x	560	0.03	F170	192.1	-
				078	103.55	104.55	sc x	1.0	15	130	230	x	6000	0.52	F1050	195.1	-
				079	104.55	105.45	sc x	0.9	45	130	515	1.0	9400	2.26	F1500	201.1	0.2
				33080	105.45	106.45	sc x	1.0	10	190	40	x	250	0.09	F45	204.1	-
																207.1	-
																210.1	-
																213.1	-
																216.1	-
																219.1	-

DEPTH (m)		ROCK DESCRIPTION	MINERALISATION	SAMPLE NO.	FROM	TO	CORE REC'D	ASSAY DATA per ppm							CORE REC'D		
FROM	TO							Sample Length	Pb	Zn	Cu	Ag - g/t	As	Se	Sn	RUN	SHORT
100.6	150.6	DACITIC LITHIC VITRIC TUFF TO LAPILLI TUFF of ACID composition		29799	100.45		Thin Section Sample - quartz-tourmaline-sulphide rock									222.1	-
100.6	103.5	As above with strong silica, carbonate and sericite alteration.	Pyrrhotite veins with associated pyrite, arsenopyrite and rare chalcopyrite forming an average of 20% of core volume.	37983	106.45	110	c	3.55	20	185	10	x	6		Ax	225.1	-
				984	110	115	c	5.0	10	150	100	1.0	4		Ax	228.1	-
				985	115	121	c	6.0	20	200	60	x	14		F390	231.1	-
103.5	105.4	Green to pale green grey crystal vitric lithic lapilli tuff of dacitic composition composed of angular pale green fragments after pumice or acid lave alongate parallel to cleavage up to 6cm long in a chloritised vitric matrix. <i>106.95 T/S. Tuff lava. andesitic (see sample) chlorite matrix</i>	Minor veins of pyrrhotite and pyrite, averaging 0.1% core volume, up to 5mm wide	29800	104.25		Thin Section Sample - andesitic? tuff-lava									234.1	-
				37910	121	122	sc	1.0	15	175	145	x	27		Fx	237.1	-
				911	122	123.6	sc	1.6	20	160	900	x	8000		F1200	240.1	-
				912	123.6	124.6	sc	1.0	25	130	5	x	23		A6	243.1	-
105.4	119.9	Pale green to green lithic vitric lappilli tuff composed of pale green fine grained angular fragments as for 103.5-105.4 and sub-angular feldspar-quartzphyric acid volcanic fragments in a chloritised vitric matrix. Calcite, epidote and chlorite veins are common (approx 10% core volume) and are up to 2cm wide.	Trace pyrite	37986	124.6	130	c	5.4	10	155	95	1.5	8		F140	246.1	-
				987	130	135	c	5.0	220	710	75	2.0	8		F340	249.1	-
				988	135	140.1	c	5.1	70	190	145	0.5	4		A92		
				37913	140.1	141.1	sc	1.0	10	125	30	x	10		A1		
				914	141.1	142.1	sc	1.0	15	105	355	x	230		F200		
				915	142.1	143.1	sc	1.0	5	85	340	x	680		F70		
119.9	121.2	Dark green grey micro gabbro dyke	Trace pyrite	916	143.1	144.1	sc	1.0	10	95	45	x	14		F15		
				917	144.1	145.1	sc	1.0	10	170	180	15	46		F130		
121.2	122	As for 105.4 to 119.9		918	145.1	146.6	sc	1.0	6	140	370	140	2.5	6		F440	
				919	146.6	147.6	sc	1.0	80	275	670	2.5	960		F920		
122	125	As above - sericitised and cleaved with fragments elongated parallel to cleavage Occasional carbonate and quartz veins (approx 5% core volume) up to 4cm wide cut this unit.	Pyrrhotite occurs as fine irregular veins and blebs representing 0.5% core volume with minor associated pyrite.	37920	147.6	148.6	sc	1.0	150	400	100	1.5	9		F740		
				921	148.6	149.6	sc	1.0	5	170	230	1.0	4		F470		
				922	149.6	150.6	sc	1.0	10	70	410	1.0	7400		F3950		
				923	150.6	151.6	sc	1.0	20	180	360	1.0	500		F2250		
				924	151.6	152.6	sc	1.0	5	120	5	0.5	20		A2		
				925	152.6	153.6	sc	1.0	15	165	275	1.0	18		A6		
125	128.9	Green lithic vitric tuff composed of pale green roughly equant angular lithic fragments in a chloritised vitric matrix.	Except for trace galena and sphalerite at 128.1 there is no visible mineralisation.	926	153.6	154.6	sc	1.0	10	260	1000	3.0	840		A4		
				927	154.6	155.6	sc	1.0	15	150	900	3.0	1.4%		Ax		
				928	155.6	156.6	sc	1.0	20	120	900	2.5	5.6%		Ax		
				929	156.6	157.6	sc	1.0	20	145	525	1.5	1.4%		Ax		
128.9	131.2	As for 119.9 to 121.2	Trace pyrite & pyrrhotite	37930	157.6	158.6	sc	1.0	5	130	210	0.5	7400		F490		
				931	158.6	159.6	sc	1.0	5	180	320	0.5	5200		F75		
131.2	141.1	Green lithic vitric tuff composed of pale green roughly equant angular lithic fragments in a chloritised vitric matrix 134.1-141.1 There is a steady increase down hole in the presence of carbonate, quartz veining and silica, chlorite alteration	<u>131.2-134.1</u> Veined pyrrhotite & pyrite up to 2cm wide making up approx 5% of 1) core volume with associated traces of chalcopyrite, galena & sphalerite.	932	159.6	160.6	sc	1.0	10	95	700	2.0	4400		F20		
				933	160.6	162.1	sc	1.0	15	110	1300	2.0	7800		F150		
				934	162.1	163.1	sc	1.0	10	220	50	0.5	180		Fx		

DEPTH (m)		ROCK DESCRIPTION	MINERALISATION	SAMPLE NO.	FROM	TO	CORE REC'D	ASSAY DATA per ppm							CORE REC'D		
FROM	TO							Sample Length	Pb	Zn	Cu	Ag - g/t	As	Fe	Sn	RUN	SHORT
			<u>134.1-141.1</u> Traces of pyrrhotite & pyrite occur as veins & blebs.														
141.1	150.6	Heavily silicified & cleaved lithic vitric tuff which is dark green grey <i>147.6 T/S altered - Agglomerate Ash flow tuff. Leuco-andesite</i>	Fine pyrrhotite, pyrite veining (average 5% core volume) with minor arsenopyrite & chalcopyrite associated with larger pyrrhotite veins near the bottom of this unit. Mineralisation localised by cleavage.	36705 706 707 708	147.4 155.1 163.1 184.4		Thin section sample - felsic intermediate tuff Thin section sample - porphyritic to glomero-porphyritic Thin section sample - porphyritic dacite Thin section sample - Quartz sericite phyllite										
150.6	163.1	DARK GREEN CLEAVED DACITIC LITHIC VITRIC TUFF composed of lithic fragments up to 10mm diameter in a vitric matrix.	<u>150.6-158.3</u> Common fine pyrrhotite veins (approx 10% core volume) with less common pyrite veins (5% core volume) both up to 1cm wide. Minor chalcopyrite is associated with pyrite along tension cracks & cleavage planes. Arsenopyrite occurs in large veins up to 3cm wide in association with larger pyrrhotite veins. There is up to 30% arsenopyrite with an average 2%	37989 990 991 992	163 165 170 175	165 170 175 179.3	c c c c	2.0 5.0 5.0 4.3	25 20 20 330	150 180 180 190	15 10 10 15	0.5 x 0.5 2.5	4 2 16 10			Ax Ax Ax A22	
150.6	152	Strong alteration (silica flooding & chloritisation) and cleavage. <i>155-1 T/S Porphyritic lava - Leuco-andesite (2 lava domes) 155-7 T/S 130-95-50 rock with silicified. porphyritic blebs.</i>	Pyrite occurs as large aggregates (up to 10% core volume) with associated less abundant pyrrhotite. Pyrrhotite also occurs as separate fine veins up to 3mm diameter.	37935 936 937 938 939 37940 941 942	179.3 180.3 181.3 182.3 183.3 184.3 185.3 186.3	180.3 181.3 182.3 183.3 184.3 185.3 186.3 187.3	sc sc sc sc sc sc sc	1.0 1.0 1.0 1.0 1.0 1.0 1.0	260 30 30 35 30 35 25	560 150 115 135 120 110 100	205 420 800 230 385 405 470	1.0 1.0 1.0 1.0 1.0 2.0 2.0	90 760 700 1120 26 38 38			F650 F450 F500 F920 F550 A6 Ax Ax	
163	178.1	Dark green fine grained possibly intrusive non-magnetic andesite, cut by carbonate and quartz veins (with carbonisation, bleaching and minor silicification from 174.0m) <i>163-1 T/S. Porphyritic lava to porphyritic andesite. (2 lava domes)</i>	Trace pyrite blebs & veins.	37993 994 995 996 997 998 999	187.3 190.0 195 200 205 210 215	190.0 195 200 205 210 215 219.1	c c c c c c c	2.7 5.0 5.0 5.0 5.0 5.0 4.1	15 20 15 5 50 10 10	100 75 55 30 200 45 55	80 130 140 30 70 25 45	1.5 1.0 1.0 x 1.0 1.5 1.0	32 40 22 14 920 20 30			Ax Ax A2 Ax Ax Ax A6	
178.1	179.3	ALTERED TUFF pale grey bleached, sheared and bedded	Trace pyrite blebs & veins.	37943 944 945 946 947	219.1 220.1 221.1 222.1 223.1	220.1 221.1 222.1 223.1 224.1	sc sc sc sc sc	1.0 1.0 1.0 1.0 1.0	15 15 25 10 10	75 70 80 60 55	100 10 205 10 15	2.0 1.0 2.5 2.0 1.0	3400 38 24 25 60	x x x x x		A8 A16 Ax Ax Ax	
179.3	249.1	"FARRELL SLATES"		948 949	224.1 225.1	225.1 226.1	sc sc	1.0 1.0	10 15	85 90	225 280	x 2.0	114 1160	0.14 0.29			Ax Ax
179.3	186.2	Heavily cleaved siltstone, greywacke & shale grey to dark grey in colour. Rare carbonate & quartz veins occur. <i>186-4 T/S 95-veins phyllite from psomopelitic gneiss</i>	Fine stringers & veins of pyrrhotite up to 30% core volume with an average of 5% with minor pyrite & trace chalcopyrite. Larger pyrrhotite veins (over 2cm wide) are associated with clear to blue	36701 37953 952 951 950 36702 703 704	226.1 231.1 232.1 233.1 234.1 235.1 240.1 245	231.1 232.1 233.1 234.1 235.1 240.1 245 249.1	c sc sc sc sc c c c	5.0 1.0 1.0 1.0 1.0 5.0 5.0 4.1	20 20 25 20 25 60 45 70	85 90 140 95 135 310 100 85	40 10 135 80 90 50 25 70	x 1.0 2.0 0.5 0.5 x x x	46 24 36 18 40 128 28 34			A3 A4 A4 Ax Ax A3 A2 A3	

ELECTROLYTIC ZINC CO OF A'ASIA LTD
ROSEBERY - TASMANIA

DIAMOND DRILL CORE RECORD

HOLE No. STP 217 4A

A 21740

DEPTH		ROCK DESCRIPTION	MINERALISATION	SAMPLE No.	8-13 FROM	14-19 TO	CORE REC'D	ASSAY DATA							CORE REC'D				
FROM	TO							Sample Length	20-25 Pb%	26-31 Zn%	32-37 Cu%	38-43 Ag - g/t	44-49 Au - g/t	50-55 Fe%	RUN	SHORT			



