

COMPANY: Golden Triangle  
 PROJECT: Main Creek Magnesite  
 HOLE NUMBER: MC 55

828241

Commenced:	05 April 99
Completed:	09 April 99
Logged By:	L.A.Newnham
Drilled By:	Almac Drilling

Purpose of Hole
To test the western half of the southern end of the Main Creek deposit at shallow depth;

Comments on Completion
intersected two high grade lenses of magnesite correlated with the D-lens (near outcrop) and D-1 lens;

Collar Details

Grid	Northing	Easting	Elevation	Dip	Bearing
AMG	5398897.3	346940.2	2097.3	-45	239

Length (m)
265.5

Hole Size	
To (m)	Size
9	HW
11.5	HQ
265.5	NQ

Significant Core Loss Zones		
From	To	%Rec.
0.0	9.0	0
251.5	253.5	65

Hole Condition on Completion
all casing and rods removed from hole; 6 m. PVC collar pipe left in hole;

Summary of Results:

Depth		Recovery	Description	Assays				
From	To			%	Length	MgO	CaO	SiO <sub>2</sub>
10.0	29.0	100	magnesite	19.0	43.87	2.97	0.52	1.51
101.0	153.0	100	magnesite	52.0	44.95	2.37	0.18	0.74
230.0	235.0	100	magnesite	5.0	41.59	2.24	3.62	3.57



# GOLDEN TRIANGLE RESOURCES N.L.

828242

A.C.N. 066 353 231

FOR AND ON BEHALF OF  
GOLDEN TRIANGLE RESOURCES N.L.  
A.C.N. 066 353 231

REGISTERED OFFICE:  
LEVEL 3  
71 QUEENS ROAD  
MELBOURNE VIC  
AUSTRALIA 3004

TEL: 61 3 9510 2544  
FAX: 61 3 9510 2770

## FACSIMILE TRANSMISSION

TO: Lindsay Newnham

FROM: Matt Noonan

DATE: 28/5/99

CC:

RECEIVER'S FAX NO:  
(03) 6394 3435

No of Pages:  
(Including this Page) (1)

RE: Grades

Lindsay,

HOLE-ID	FROM	TO	INTERVAL	MGO	CAO	FE2O3	SIO2
MC 55	10	29	19	43.87	2.97	1.51	0.52
MC 55	101	153	52	44.95	2.37	0.74	0.18
MC 55	230	235	5	41.59	2.24	3.57	3.62
MC 56	134	193	59	44.07	2.00	0.72	2.82

Matt



828243

**GOLDEN TRIANGLE RESOURCES N.L.**

A.C.N. 066 353 231

FOR AND ON BEHALF OF  
GOLDEN TRIANGLE RESOURCES N.L.  
A.C.N. 066 353 231

REGISTERED OFFICE:  
LEVEL 3  
71 QUEENS ROAD  
MELBOURNE VIC  
AUSTRALIA 3004

**FACSIMILE TRANSMISSION**

TEL: 61 3 9510 2544  
FAX: 61 3 9510 2770

TO: Lindsay Newnham

FROM: Matt Noonan

DATE: 19/5/99

CC:

RECEIVER'S FAX NO:

(03) 6394 3435

No of Pages:

(Including this Page) (1)

RE: Grades

Lindsay,



Results as Requested

HOLE-ID	FROM	TO	INTERVAL	CAO	FE2O3	MGO	SiO2
MC 30	48.4	119.1	69.7	2.51	1.50	43.93	1.31
MC 30	133	151	16.6	2.41	0.66	45.44	0.16
MC 30	161.9	174	12	2.44	0.40	45.21	0.10
MC 30	185	199	14	2.50	0.68	45.05	0.10
MC 34	195.3	201.5	6.2	1.98	2.76	43.49	1.67
MC 34	209.3	217.3	8	2.82	2.72	41.94	3.41
MC 34	241.7	246.7	5	2.07	2.44	40.44	8.45
MC 34	297.1	306.1	9	2.09	1.74	41.93	5.51
MC 34	441.1	446.1	5	2.55	0.83	44.16	1.56
MC 36	171.2	212.2	41	2.44	0.78	43.69	2.50
MC 53	23	52	29	2.01	3.98	42.80	0.63
MC 53	64	85	21	1.57	5.52	41.23	1.87
MC 53	139	160	21	2.70	3.07	42.87	0.77
MC 53	180.5	219	37.2	2.23	1.06	41.54	7.62
MC 53	263	302	33	2.53	0.94	44.34	0.96
MC 53	330.5	369	37	2.34	0.63	44.95	0.38
MC 53	397	404	7	1.78	0.72	45.33	0.26
MC 55	10	42	28.7	3.19	1.65	43.24	1.20
MC 55	101	152	51	2.36	0.73	44.96	0.18
MC 56	56	72	16	2.03	1.80	42.94	3.41

Matt



828245



5 cm

MC55

COMPANY: Golden Triangle NL  
PROJECT: Main Creek  
HOLE NUMBER: MC 55

Page No: 1

Description		Core Recovery			RQD			Assays									
From	To		From	To	%	From	To	%	From	To	MgO	CaO	SiO <sub>2</sub>	Fe <sub>2</sub> O <sub>3</sub>			
0.0	9.0	<b>HW TRICONE, no core:</b>	0.0	9.0	0												
9.0	34.7	<b>MAGNESITE:</b> white magnesite, extensively replaced by clear-white crystalline magnesite; coarse crystalline magnesite present as 1-50 mm. wide veins and irregular masses; becomes slightly grayer (dolomitic) towards base of interval; occasional small seggregations gray quartz; no talc observed; very rare fine grained pyrite associated with crystalline magnesite; core fresh but moderately broken due to joint sets at 10°, 30°, 50° CA combining with fracturing along thin crystalline magnesite veins; some joint surfaces 15-22.5 m. limonite stained and possibly transmitting water;	9.0	34.7	100	9.0	11.5	60	9.0	10.0	41.38	6.16	0.65	1.41			
						11.5	15.4	35	10.0	11.0	43.20	3.63	1.05	1.48			
						15.4	19.5	50	11.0	12.0	44.48	2.24	0.62	1.23			
						19.5	23.9	70	12.0	13.0	44.29	2.81	<0.05	1.17			
						23.9	28.3	75	13.0	14.0	45.04	1.60	1.29	1.28			
						28.3	32.8	85	14.0	15.0	44.56	2.04	0.77	1.29			
						32.8	37.5	75	15.0	16.0	43.87	3.04	0.84	1.37			
									16.0	17.0	43.17	3.80	0.42	1.56			
									17.0	18.0	43.34	3.61	0.62	1.46			
									18.0	19.0	43.18	2.91	2.05	1.46			
									19.0	20.0	44.02	2.64	0.45	1.68			
									20.0	21.0	44.28	2.66	<0.05	1.64			
									21.0	22.0	44.48	2.75	<0.05	1.50			
									22.0	23.0	44.16	3.05	<0.05	1.61			
									23.0	24.0	43.26	3.36	0.17	1.59			
									24.0	25.0	43.07	3.68	0.75	1.67			
									25.0	26.0	42.68	3.76	0.27	1.75			
34.7	38.0	<b>INTERBEDDED SCHIST and MAGNESITE:</b> light-dark gray schist, variably talcose and with abundant carbonate spotting in places; SCA 45-50°; <b>34.7-35.3 m:</b> schist with some magnesite bands; <b>35.3-36.9 m:</b> magnesite replaced by crystalline magnesite; dolomitic in part; pug seam on FW; <b>36.9-38.0 m:</b> dark schist, strongly carbonate spotted below 37.4 m; core moderately competent but fractures parallel to schistosity common in schist;	34.7	36.9	90				26.0	27.0	42.74	4.07	0.47	1.83			
			36.9	38.0	100				27.0	28.0	44.61	2.69	<0.05	1.58			
									28.0	29.0	45.05	2.15	<0.05	1.47			
									29.0	30.0	43.86	3.25	<0.05	1.66			
									30.0	31.0	41.74	4.80	0.10	1.72			
									31.0	32.0	42.66	4.31	0.42	1.66			
									32.0	33.0	42.41	4.31	0.22	1.99			
									33.0	34.7	42.21	3.70	0.72	2.67			
									38.0	39.0	39.92	4.97	5.80	1.72			
									39.0	40.0	41.62	2.52	5.20	1.77			
									40.0	41.0	41.96	2.14	5.76	1.85			
									41.0	42.0	41.48	2.38	5.74	1.84			
38.0	52.7	<b>MAGNESITE:</b> white magnesite, extensively replaced by clear-light gray (dolomitic ?) crystalline magnesite; abundant thin veins of coarse crystalline magnesite; overall wispy appearance; small irregular patches of light gray silica common; no talc observed; minor fine grained disseminated pyrite .....	38.0	52.7	100	37.5	42.2	95	42.0	43.0	35.60	7.02	9.55	2.01			
						42.2	46.7	90	43.0	44.0	34.82	6.59	11.64	2.43			
						46.7	51.2	100	44.0	45.0	39.69	4.68	5.22	2.08			
									45.0	46.0	40.56	3.52	5.80	1.94			
									46.0	47.0	36.37	7.62	7.38	2.14			
									47.0	48.0	37.46	6.46	6.58	2.17			
									48.0	49.0	37.39	5.41	8.46	2.22			
									49.0	50.0	41.06	3.71	5.13	1.49			

COMPANY: Golden Triangle NL  
 PROJECT: Main Creek  
 HOLE NUMBER: MC 55

Page No: 2

Description		Core Recovery			RQD			Assays								
From	To		From	To	%	From	To	%	From	To	MgO	CaO	SiO <sub>2</sub>	Fe <sub>2</sub> O <sub>3</sub>		
38.0	52.7	associated with replacement; <b>below 51.0 m:</b> core surface of magnesite is creamy color (ie) dolomitic magnesite; principal joint set 30° CA; in general, ground conditions excellent; grades into.....							50.0	51.0	40.81	1.94	8.03	1.66		
continued.....									51.0	52.7	29.34	12.46	14.47	1.20		
52.7	56.8	<b>MAGNESITE, dolomitic, silicified, pyritic:</b> whitemagnesite almost completely replaced by strongly silicified light-medium gray dolomite; in places core is totally white fractured quartz; no talc observed; 1-2% pyrite associated with dolomite and quartz; locally 3-4%; ground conditions excellent;	52.7	56.8	100	51.2	55.8	100								
56.8	89.8	<b>INTERBEDDED SCHIST and MAGNESITE:</b> <b>56.8 m:</b> 200 mm light gray broken schist; <b>56.8-57.1 m:</b> white magnesite; <b>57.1-61.3 m:</b> dark gray, carbonate spotted, schistose volcanic, with 200 mm magnesite band at 59.5 m; crumbly sericitic schist band at base; SCA 50-60°; <b>61.3-62.9 m:</b> white magnesite extensively replaced by crystalline magnesite; 1-10 mm coarse crystalline magnesite veins common; good ground conditions; <b>62.9-69.0 m:</b> dark gray, fine grained schistose volcanic; numerous microveins of white carbonate; occasional veins up to 30 mm. of coarse crystalline magnesite; weakly magnetic in places; SCA variable but generally 60°; basal 400 mm. strongly talcose; <b>69.0-76.0 m:</b> white magnesite extensively replaced by tight gray (? dolomitic) crystalline magnesite; irregular patches of gray silica common; talc is associated with alteration but overall is a minor component; minor disseminated fine grained pyrite associated with crystalline magnesite, talc and quartz alteration; HW 200 mm., and FW 100 mm. strongly talcose;	56.8	89.8	100	55.8	60.3	35	69.0	70.0	32.23	10.94	11.79	3.14		
						60.3	64.6	80	70.0	71.0	36.40	7.08	9.44	2.85		
						64.6	69.4	65	71.0	72.0	40.90	3.97	4.54	2.67		
						69.4	74.0	100	72.0	73.0	39.66	4.61	5.98	2.37		
						74.0	78.3	90	73.0	74.0	39.13	6.11	5.12	2.02		
						78.3	83.0	75	74.0	75.0	38.48	5.79	5.34	2.20		
						83.0	87.5	80	75.0	76.0	37.57	8.07	4.07	2.37		
						87.5	92.0	50								

COMPANY: Golden Triangle NL  
PROJECT: Main Creek  
HOLE NUMBER: MC 55

Page No: 3

Description		Core Recovery			RQD			Assays							
From	To		From	To	%	From	To	%	From	To	MgO	CaO	SiO <sub>2</sub>	Fe <sub>2</sub> O <sub>3</sub>	
56.8 continued.....	89.8	principal joint set 50° CA; ground conditions very good except for talcose zone on HW; <b>76.0-79.3 m:</b> dark gray fine grained schistose volcanic; talcose and strongly altered near margins; SCA 35°-40°; ground conditions good; <b>79.3-79.9 m:</b> cream and gray banded carbonate; <b>79.7-80.7 m:</b> dark and fine grained schistose volcanic; <b>80.7-82.6 m:</b> magnesite, extensively replaced by talc, dolomite and quartz; minor 1-5 mm crystalline magnesite veining; <b>82.6-86.3 m:</b> dark gray fine grained schistose volcanic; numerous narrow disrupted quartz and quartz carbonate veins; talcose puggy band on FW; <b>86.3-87.8 m:</b> white magnesite, extensively replaced; <b>87.8-89.8 m:</b> dark gray-green weakly schistose volcanic; carbonate veining common; SCA 60°; ground reasonably competent but some broken sections;													
89.8	154.5	<b>MAGNESITE:</b> white magnesite, variably mottled and veined by coarse crystalline magnesite; <b>89.8-105.0 m:</b> white magnesite, fragmented and extensively replaced by light gray crystalline magnesite resulting in mottled texture; possibly dolomitic in places; numerous 1-10 mm. veins of coarse crystalline magnesite; only trace talc; minor fine grained disseminated pyrite associated with crystalline magnesite; principal joint direction 30° CA; ground conditions excellent; grades into..... <b>105.0-150.5 m:</b> similar to magnesite above but less of a mottled texture (ie) less crystalline magnesite and possibly less dolomitic; greater abundance of coarse.....	89.8	154.5	100	92.0	96.6	90	90.0	91.0	43.33	3.87	0.19	1.07	
						96.6	101.0	100	91.0	92.0	43.45	3.80	0.21	0.90	
						101.0	105.6	85	92.0	93.0	44.77	2.78	<0.05	0.83	
						105.6	110.2	95	93.0	94.0	44.01	3.44	<0.05	0.91	
						110.2	114.8	100	94.0	95.0	41.71	5.47	<0.05	0.80	
						114.8	119.4	95	95.0	96.0	42.54	5.21	<0.05	0.81	
						119.4	123.5	80	96.0	97.0	36.43	11.84	0.31	1.23	
						123.5	128.2	100	97.0	98.0	38.04	10.09	0.22	1.09	
						128.2	132.6	80	98.0	99.0	43.36	4.19	0.17	0.80	
						132.6	137.1	95	99.0	100.0	43.16	4.38	0.10	0.88	
						137.1	141.5	85	100.0	101.0	44.16	3.43	<0.05	0.72	
						141.5	146.2	95	101.0	102.0	45.21	2.09	<0.05	0.64	
						146.2	151.0	95	102.0	103.0	45.41	2.25	<0.05	0.64	
						151.0	155.6	95	103.0	104.0	45.21	2.20	<0.05	0.69	
									104.0	105.0	45.96	1.72	<0.05	0.66	
									105.0	106.0	45.96	1.72	<0.05	0.60	

COMPANY: Golden Triangle NL  
PROJECT: Main Creek  
HOLE NUMBER: MC 55

Page No: 4

Description		Core Recovery			RQD			Assays								
From	To		From	To	%	From	To	%	From	To	MgO	CaO	SiO <sub>2</sub>	Fe <sub>2</sub> O <sub>3</sub>		
89.8	154.5	crystalline magnesite 1-10 mm veins; very rare fine grained pyrite associated with crystalline magnesite; no talc observed; principal joint set widely spaced 30° CA; ground conditions excellent; many breaks are driller breaks; grades into..... <b>150.5-154.5 m:</b> similar to unit above but pervasive pale gray color (? more dolomitic); abundant coarse crystalline magnesite veining constituting up to 50% of core in places; no talc observed; very rare fine grained pyrite; ground conditions excellent; grades into.....							106.0	107.0	45.85	1.66	<0.05	0.66		
									107.0	108.0	45.24	2.18	<0.05	0.65		
									108.0	109.0	45.45	1.72	0.15	0.81		
									109.0	110.0	45.83	0.93	0.87	0.73		
									110.0	111.0	43.32	4.00	0.44	0.70		
									111.0	112.0	42.87	5.03	0.23	0.79		
									112.0	113.0	45.43	1.79	0.29	0.65		
									113.0	114.0	46.31	0.85	<0.05	0.81		
									114.0	115.0	44.01	3.42	0.38	0.83		
									115.0	116.0	43.61	4.05	0.19	0.66		
									116.0	117.0	44.84	2.29	0.40	0.72		
									117.0	118.0	44.38	2.47	<0.05	0.63		
									118.0	119.0	45.30	1.77	<0.05	0.73		
									119.0	120.0	44.93	2.17	0.14	0.77		
154.5	156.1	<b>SCHIST:</b> dark gray-green fine grained weakly schistose volcanic with talcose margins; minor fine grained pyrite; ground moderately competent except for weak talcose margins;	154.5	156.1	100				120.0	121.0	46.39	0.66	<0.05	0.95		
									121.0	122.0	46.47	0.68	<0.05	0.64		
									122.0	123.0	46.72	0.16	0.17	0.78		
									123.0	124.0	46.41	0.39	0.25	0.92		
									124.0	125.0	45.95	1.02	0.14	0.93		
									125.0	126.0	45.11	2.62	<0.05	0.60		
									126.0	127.0	43.56	4.10	0.19	0.78		
156.1	165.0	<b>MAGNESITE:</b> <b>156.1-162.2 m:</b> white magnesite extensively brecciated and replaced by dolomitic crystalline magnesite, resulting in a strong mottled appearance; minor fine grained disseminated pyrite associated with dolomite-crystalline magnesite; <b>162.2-165.0 m:</b> white magnesite extensively brecciated and replaced by light gray-white crystalline magnesite; abundant 1-10 mm veins coarse crystalline magnesite; very rare fine grained pyrite; core very competent; not assayed because of dolomitic upper section;	156.1	165.0	100	155.6	160.2	95	127.0	128.0	45.07	2.02	0.53	1.05		
						160.2	164.9	95	128.0	129.0	43.14	4.01	0.26	0.74		
									129.0	130.0	43.77	4.07	0.20	0.64		
									130.0	131.0	45.43	1.59	0.22	0.65		
									131.0	132.0	45.82	1.63	<0.05	0.61		
									132.0	133.0	45.39	1.82	0.25	0.68		
									133.0	134.0	45.63	1.65	<0.05	0.74		
									134.0	135.0	44.77	2.78	0.19	0.67		
									135.0	136.0	44.93	2.29	0.26	0.65		
									136.0	137.0	45.11	2.04	0.39	0.72		
									137.0	138.0	45.49	1.91	0.23	0.69		
									138.0	139.0	44.52	2.74	<0.05	0.71		
									139.0	140.0	43.88	4.02	<0.05	0.74		
									140.0	141.0	43.64	3.65	0.27	0.65		
									141.0	142.0	44.81	2.72	0.16	0.72		
165.0	171.3	<b>SCHIST:</b> dark gray-black schistose ? sediment; strongly talcose; bedding parallels schistosity 40° CA; possibly a sheared shale or siltstone; <b>below 170.5 m:</b> black, carbonaceous, pyritic;	165.0	171.3	100	164.9	169.5	55	142.0	143.0	45.07	2.46	0.24	0.67		
						169.5	174.0	75	143.0	144.0	44.40	3.05	0.50	0.68		
									144.0	145.0	44.49	3.08	0.35	0.71		
									145.0	146.0	44.58	2.57	0.17	0.69		

COMPANY: Golden Triangle NL  
PROJECT: Main Creek  
HOLE NUMBER: MC 55

Page No: 5

Description		Core Recovery			RQD			Assays										
From	To		From	To	%	From	To	%	From	To	MgO	CaO	SiO <sub>2</sub>	Fe <sub>2</sub> O <sub>3</sub>				
165.0	171.3	core above 170.5 m., soft and broken along schistosity/bedding; 150 mm pug zone on HW; below 170.5 m., core very broken; irregular FW contact;							146.0	147.0	44.53	2.67	<0.05	0.70				
continued.....									147.0	148.0	44.40	3.04	0.10	0.73				
										148.0	149.0	45.39	2.20	<0.05	0.74			
										149.0	150.0	44.75	2.79	<0.05	0.82			
										150.0	151.0	44.40	2.74	0.16	0.88			
171.3	205.3	<b>MAGNESITE, dolomitic:</b> mottled dolomitic magnesite with increasing pyrite towards base of interval; <b>171.3-172.1 m:</b> dark gray dolomite; <b>172.1-189.8 m:</b> white magnesite extensively brecciated and replaced by light gray dolomitic crystalline magnesite, resulting in a pale mottled appearance; 1-5 mm veins of coarse crystalline magnesite common; 20 mm. talcose schist band at 180.5 m; minor fine grained disseminated pyrite associated with dolomitic crystalline magnesite and also along late stage fine fractures; <b>189.8-190.2 m:</b> soft dark gray talcose schist; <b>190.2-194.5 m:</b> magnesite similar to 172.1 m....., but increasing pyrite towards base (still <1%), accompanied by patches of talc; <b>194.5-194.8 m:</b> soft dark brown talcose schist; SCA 70°; <b>194.8-205.3 m:</b> white magnesite extensively replaced by light gray dolomitic crystalline magnesite; abundant 1-10 mm veins of coarse crystalline magnesite; minor talc associated with replacement; 1-2% fine to coarse disseminated pyrite, gradually increasing down hole; below 202 m., 3-5% pyrite as disseminations and infilling stylolitic structures; soft talcose schist partings towards base of interval; overall ground conditions very good except for schist zones which are soft and broken;	171.3	205.3	100	174.0	178.6	90	151.0	152.0	43.94	3.00	0.35	0.89				
										178.6	183.2	95	152.0	153.0	44.10	2.97	0.19	1.11
										183.2	187.6	90	153.0	154.5	40.56	6.85	0.74	1.26
										187.6	192.3	95						
										192.3	196.9	90	173.0	174.0	43.16	4.29	<0.05	0.78
										196.9	201.4	100	174.0	175.0	41.82	5.88	<0.05	0.88
										201.4	206.1	85	175.0	176.0	42.48	5.50	<0.05	0.76
										176.0	177.0	42.34	5.08	0.56	0.70			
										177.0	178.0	42.35	5.80	0.26	0.73			
										178.0	179.0	43.19	4.26	0.87	0.74			
										179.0	180.0	42.50	5.08	0.26	0.87			
										180.0	181.0	40.21	6.84	1.64	1.05			
										181.0	182.0	41.24	6.28	0.83	0.90			
										182.0	183.0	40.36	7.83	0.48	0.85			
										183.0	184.0	43.47	3.93	0.23	0.80			
									184.0	185.0	42.39	4.92	0.28	0.85				
									185.0	186.0	40.84	6.61	0.50	0.97				
									186.0	187.0	40.78	7.07	<0.05	0.99				
									187.0	188.0	39.46	8.75	0.18	1.20				
									188.0	189.0	38.25	9.33	0.56	1.44				
									189.0	190.0	39.18	7.53	1.85	1.54				
									190.0	191.0	39.67	7.53	1.05	1.70				
									191.0	192.0	40.68	6.22	0.78	2.24				
									192.0	193.0	38.08	9.08	0.46	2.35				
									193.0	194.0	37.00	10.02	2.13	1.97				
									194.0	195.0	38.39	7.75	2.58	2.44				
									195.0	196.0	40.76	6.50	<0.05	2.04				
									196.0	197.0	39.79	7.39	1.78	1.46				
									197.0	198.0	30.62	18.65	0.80	1.51				
									198.0	199.0	26.37	23.01	2.56	1.36				
									199.0	200.0	29.96	18.46	2.48	1.17				
									200.0	201.0	31.73	16.87	1.41	1.72				
205.3	207.0	<b>SCHIST:</b> dark gray talcose pyritic schist; strongly pyritic with large aggregates and semi-massive	205.3	207.0	100	206.1	210.6	70										

COMPANY: Golden Triangle NL  
PROJECT: Main Creek  
HOLE NUMBER: MC55

Page No: 6

Description		Core Recovery			RQD			Assays								
From	To		From	To	%	From	To	%	From	To	MgO	CaO	SiO <sub>2</sub>	Fe <sub>2</sub> O <sub>3</sub>		
205.3 continued.....	207.0	sulfide bands in places; SCA 40°; core very soft and broken, mainly along schistosity; soft puggy seam on FW;														
207.0	249.5	<b>MAGNESITE, dolomitic and pyritic:</b> magnesite, variably altered and replaced; <b>207.0-208.0 m:</b> intensely altered magnesite with common talc and 1% disseminated pyrite; thin discontinuous schist partings; <b>208.0-212.0 m:</b> massive white magnesite; coarse crystalline magnesite common as 1-10 mm. veins; ground conditions excellent; grades into..... <b>212.0-219.0 m:</b> white chalky magnesite extensively brecciated and replaced by light gray dolomitic crystalline magnesite; evidence of minor water leaching (ie) vuggy; 216.3 m: 300 mm band 2-3% coarse disseminated pyrite; grades into..... <b>219.0-229.0 m:</b> mottled dolomitic unit with white magnesite brecciated and replaced by light gray dolomite and crystalline magnesite;; 1-2% pyrite as disseminated grains associated with alteration, and as thin seams along fractures and in stylolites; ground conditions excellent; grades into..... <b>229.0-236.0 m:</b> white magnesite extensively altered and replaced by abundant crystalline and coarse crystalline magnesite; minor talc component; minor medium grained pyrite, often >1% in narrow bands; principal joint direction 30° CA; ground conditions excellent; grades into..... <b>236.0-240.0 m:</b> massive white magnesite, remaining as large ghost like lumps, almost totally replaced by light gray dolomitic crystalline magnesite; coarse crystalline magnesite common as 1-5 mm veins; minor talc patches; trace fine grained disseminated pyrite; ground conditions excellent; grades into.....	207.0	249.5	100	210.6	215.3	100	208.0	209.0	40.49	6.28	0.87	2.20		
						215.3	220.0	95	209.0	210.0	43.78	2.55	<0.05	1.62		
						220.0	224.7	100	210.0	211.0	44.36	2.84	<0.05	1.08		
						224.7	229.4	95	211.0	212.0	43.57	3.69	<0.05	0.93		
						229.4	233.9	85	212.0	213.0	33.86	14.88	0.36	1.27		
						233.9	238.3	95	213.0	214.0	41.08	7.32	<0.05	0.70		
						238.3	242.8	95	214.0	215.0	39.52	8.45	0.15	1.26		
						242.8	247.8	90	215.0	216.0	42.40	5.09	0.12	0.83		
						247.8	252.5	85	216.0	217.0	34.69	13.24	0.34	1.62		
									217.0	218.0	33.31	14.81	0.90	1.28		
									218.0	219.0	40.13	7.95	<0.05	1.06		
									229.0	230.0	40.88	4.47	0.26	3.61		
									230.0	231.0	41.81	2.26	3.25	3.45		
									231.0	232.0	40.59	2.08	5.56	3.97		
									232.0	233.0	40.91	2.76	2.82	3.77		
									233.0	234.0	42.75	1.90	2.20	3.33		
									234.0	235.0	41.91	2.22	4.29	3.33		
									235.0	236.0	41.01	3.70	3.45	2.52		
									236.0	237.0	42.58	2.92	3.41	1.59		
									237.0	238.0	42.92	3.38	1.20	1.26		
									238.0	239.0	43.32	3.40	<0.05	1.37		
									239.0	240.0	39.88	8.09	<0.05	0.95		
									240.0	241.0	38.35	8.13	3.54	1.53		
									241.0	242.0	33.99	8.97	12.68	1.69		
									242.0	243.0	39.49	4.17	13.84	0.97		
									243.0	244.0	40.14	3.42	12.60	1.18		
									244.0	245.0	43.43	1.94	6.11	1.00		
									245.0	246.0	40.69	3.71	8.39	1.15		
									246.0	247.0	31.88	14.03	8.60	1.61		
									247.0	248.0	36.54	10.60	1.62	2.30		
									248.0	249.5	32.30	12.46	6.57	2.58		

COMPANY: Golden Triangle NL  
 PROJECT: Main Creek  
 HOLE NUMBER: MC 55

Page No: 7

Description		Core Recovery			RQD			Assays										
From	To		From	To	%	From	To	%	From	To	MgO	CaO	SiO <sub>2</sub>	Fe <sub>2</sub> O <sub>3</sub>				
207.0 continued.....	249.5	<b>240.0-249.5 m:</b> white magnesite, extensively altered and replaced by white and green talc, accompanied by dolomitic magnesite and free irregular patches of silica; minor fine-medium grained euhedral disseminated pyrite associated with alteration zones; ground competent but weak in more talcose sections;																
249.5	251.5	<b>INTERBEDDED DOLOMITE and MAGNESITE:</b> white magnesite extensively replaced by gray dolomitic crystalline magnesite, interbedded with medium gray dolomite, strongly stylolitic; 2-3% pyrite as medium-coarse disseminated grains and concentrated along fractures and stylolitic structures; 251.4 m: 15-20% semi-massive pyrite band; BCA (?) 60°;	249.5	251.5	100													
251.5	253.5	<b>ALTERED CARBONATE:</b> <b>251.5-252.1 m:</b> hematitic carbonate, largely altered to talc and quartz, containing abundant coarse black magnetite; distinctive reddish coloration; contains a 50 mm. talcose schist band; <b>252.1-252.8 m:</b> light gray dolomitic magnesite; <b>252.8-253.5 m:</b> cavity;	251.5	253.5	65	252.5	257.6	60 (+ cavity)										
253.5	256.6	<b>SCHIST:</b> dark gray-green weakly schistose volcanic with white flecking due to carbonate alteration; SCA 70°; broken pyritic HW; soft talcose FW; apart from HW and FW zones, ground conditions good with most fractures parallel to schistosity;	253.5	256.6	100	257.6	262.5	65										

